An antidepressant

Fluvoxamine (flu-vox-a-meen)

You should read this leaflet with your carer/care giver.




Fluvoxamine:

- Is also called Faverin or Luvox.
- Come as tablets or capsules.
- Fluvoxamine is used for depression, anxiety, phobias and OCD (obsessive compulsive disorder).
- Fluvoxamine is sometimes used for people who are distressed. There is no clear evidence that it improves someone's behaviour.



The label on your fluvoxamine packet should tell you when and how much medicine to take.

Swallow your tablet or capsule with water.





You should carry on taking your medicine even if you feel better unless your doctor asks you to stop.

Drinking alcohol when taking fluvoxamine could make you feel sleepy.

Some medicines do not work well together.

You should tell your doctor about any other medicines you are taking.

This is an unofficial leaflet produced by Professor Shoumitro (Shoumi) Deb and Ms. Bharati Limbu of Imperial College London, UK and Dr Gemma Unwin, University of Birmingham, UK with help from Cornwall Learning Disability Advisory Group. The project is funded by the National Institute for Health Research (NIHR), UK and sponsored by ©CNWL NHS Trust UK. Any adaptation or translation of this leaflet has to acknowledge fully the original source with author affiliation and copyright information and be made available free of charge. 01.07.2020.

Side effects:

- Medicine can have some effects on your body that are not wanted.
- Not everyone taking a medicine will get them.
- Many will go away with time.
- Many are rare.

You might want to talk to your doctor or the pharmacist or your key worker if you have any side effect.

These are some (but NOT all) side effects of fluvoxamine...



- Feeling sick or being sick
- Feeling more anxious or on edge
- Lose weight or may put on weight (check weight regularlt)
- · Problem sleeping
- Having the 'runs' or needing the toilet a lot
- Dry mouth

Headache

- Not feeling hungry
- Loss of interest in sex
- Irregular heartbeat (check ECG)



Some side effects are more serious.

You should tell someone straight away if you have...

- Thoughts about harming or killing yourself
- Fit or seizure
- Elevated mood or feel angry
- Red rashes
- Shaky or stiff
- High temperature



You should talk to your doctor before you stop taking your medicine.

Sometimes, if you stop taking your medicine, you may have:

- Flu-like symptoms
- Dizziness
- · Anxiety or feel on edge
- Problems sleeping
- Problems with memory
- Movement disorder



This is an unofficial leaflet produced by Professor Shoumitro (Shoumi) Deb and Ms. Bharati Limbu of Imperial College London, UK and Dr Gemma Unwin, University of Birmingham, UK with help from Cornwall Learning Disability Advisory Group. The project is funded by the National Institute for Health Research (NIHR), UK and sponsored by ©CNWL NHS Trust UK. Any adaptation or translation of this leaflet has to acknowledge fully the original source with author affiliation and copyright information and be made available free of charge. 01.07.2020.



This leaflet does not tell you everything about fluvoxamine.

You can get more information from:

- your pharmacist, doctor, nurse or carer/ care giver
- check
 http://www.choiceandmedication.org/cnwl/
- NHS on 111
- medicine information helpline on 020 3317 5090.

This leaflet can be downloaded free of charge from:

https://spectrom.wixsite.com/project/

While preparing this leaflet we checked information in...

- The British National Formulary (<u>www.bnf.org</u>)
- Dulcan, M. K., & Ballard, R. (2015). Helping parents and teachers understand medications for behavioural and emotional problems: A resource book of medication information handouts. American Psychiatric Publishing, Washington DC, USA.

This leaflet is not designed for people with intellectual/learning disabilities to read on their own. Support (care) staff or care givers should go through the leaflet with the person who has intellectual disabilities.

All images copyright @Photosymbols.

This is an unofficial leaflet produced by Professor Shoumitro (Shoumi) Deb and Ms. Bharati Limbu of Imperial College London, UK and Dr Gemma Unwin, University of Birmingham, UK with help from Cornwall Learning Disability Advisory Group. The project is funded by the National Institute for Health Research (NIHR), UK and sponsored by ©CNWL NHS Trust UK. Any adaptation or translation of this leaflet has to acknowledge fully the original source with author affiliation and copyright information and be made available free of charge. 01.07.2020.